



**Go in stronger. Get home sooner.**



**Reduce risk of complications in major elective surgery with IMPACT Advanced Recovery® Drink.**

- ✔ Successfully used in conjunction with Enhanced Recovery Protocols (ERPs) to reduce infection,<sup>1,2</sup> length of stay,<sup>1,3,4</sup> readmission<sup>5</sup> and hospital cost<sup>4</sup>
- ✔ Contains a proven blend of 3 specialized immunonutrients: arginine, omega-3 fatty acids and dietary nucleotides supported by the American Society of Enhanced Recovery (ASER) Implementation Guide<sup>6,7\*</sup>
- ✔ Meets the Perioperative Quality Initiative (POQI) Joint Consensus recommendations for perioperative immunonutrition.<sup>8</sup>
- ✔ Studied to help reduce hospital acquired infections when provided as part of a preoperative patient kit.<sup>9</sup>
- ✔ Accommodates carbohydrate controlled diets (20 g carbohydrate/serving)

**MAJOR SURGERY PLACES UNIQUE NUTRITIONAL DEMANDS ON THE BODY.**

IMPACT® formulas are designed with a unique blend of nutrients to meet the distinct nutritional needs of major surgery patients. They have been shown to support the body's immune response, reducing the risk of infectious complications in a variety of major surgeries,<sup>1-3, 5, 7, 10-12</sup> including:

- Gastrointestinal Cancer
- Head and Neck Cancer
- Cardiac
- Gynecologic Oncology
- Bladder Cancer
- Ventral Hernia Repair
- Orthopedic

**For more information and IMPACT® protocol resources, call 1-800-422-2752 or visit: [www.NestleMedicalHub.com](http://www.NestleMedicalHub.com).**

References: **1.** Goncalves TJM et al. *JPEN* 2020 on-line first; 0:1-8. **2.** Hamilton-Reeves JM, et al. *Euro Urol* 2016;69(3):389-392. **3.** Alito Aprelino M and de Aguilar-Nascimento JE. *Nutr J* 2016;15:1-7. **4.** Robinson LA et al. *J Thorac Cardiovasc Surg* 2020; on-line ahead of print. **5.** Majumder A, et al. *J Am Coll Surg* 2016;222:1106-1115. **6.** ASER Enhanced Recovery Implementation Guide. Sept 2016. <http://aserhg.org>. **7.** Adiamah A et al. *Ann Surg* 2019; 270(2):247-256. **8.** Wischmeyer PE et al. *Anesth Analg* 2018; 126(6):1883-1895. **9.** Kelley KE et al. *Ann Surg* 2018; 268(4):650-656. **10.** Drover JW, et al. *J Am Coll Surg* 2011;212(3):385-399. **11.** Braga M, et al. *Surg* 2002;132:805-814. **12.** Chapman JS et al. *Obstet Gynecol* 2016;128:138-144.

**USE UNDER MEDICAL SUPERVISION. NOT INDICATED WHEN IMMUNE SUPPRESSION IS DESIRED.**

\*The above statement does not constitute an endorsement of IMPACT formula or any other Nestlé Health Science product by the American Society of Enhanced Recovery (ASER).

